

EEB 581, Problem Set Four Solutions

1 : Consider a one-way ANOVA design with 5 factors and 10 replicates per factor. Suppose that factor variance σ_τ^2 is ten percent of the total variance σ_T^2 (i.e., $\sigma_\tau^2/\sigma_T^2 = 0.10$).

- (a) Given that the total variance equals the treatment plus error variance ($\sigma_T^2 = \sigma_\tau^2 + \sigma_e^2$), what is σ_τ^2/σ_e^2 ?

Note that $\sigma_T^2 = \sigma_\tau^2 + \sigma_e^2$ implies $\sigma_T^2/\sigma_\tau^2 = 1 + \sigma_e^2/\sigma_\tau^2$, or that $\sigma_e^2/\sigma_\tau^2 = \sigma_T^2/\sigma_\tau^2 - 1$, and hence

$$\sigma_\tau^2/\sigma_e^2 = \frac{1}{\sigma_T^2/\sigma_\tau^2 - 1} = \frac{1}{1/0.1 - 1} = \frac{1}{9} = 0.1111$$

- (b) What is the 95% critical value for the F-test?

The 95% critical value f satisfies $\Pr(F_{N-1, N(n-1)} \leq f) = 0.95$. Here $N = 5, n = 10$. Using R, we find that

```
> qf(0.95, 5-1, 5*(10-1))
[1] 2.578739
```

- (c) What is the power of this design (assuming a test of $\alpha = 0.05$) for a fixed-effects ANOVA?

From Equation A5.23 in the notes, the power is given by

$$\Pr(F_{N-1, N(n-1), \lambda} \geq 2.578739)$$

where the noncentrality parameter $\lambda = n(N-1)\sigma_\tau^2/\sigma_e^2 = 10 * 4 * (1/9)$. In R, we can compute this by using

```
> 1-pf(2.578739, 5-1, 5*(10-1), 10*4/9)
[1] 0.3207070
```

Note that we used 1-pf, as pf returns the probability of being $\leq x$, while we wish the probability of being $\geq x$. Note we could also have used

```
> 1-pf(qf(0.95, 5-1, 5*(10-1)), 5-1, 5*(10-1), 10*4/9)
```

directly having R recompute the critical value.

- (d) What is the power of this design under a random-effects ANOVA?

Recalling Equation A5.30b in the notes, the power is given by

$$\Pr \left[F_{N-1, N(n-1)} > \frac{F_{N-1, N(n-1), [1-\alpha]}}{1 + n(\sigma_\tau^2/\sigma_e^2)} \right] = \Pr \left[F_{N-1, N(n-1)} > \frac{2.578739}{1 + n(\sigma_\tau^2/\sigma_e^2)} \right]$$

In R,

```
> 1- pf(2.578739/(1+10/9), 5-1, 5*(10-1))
[1] 0.3150735
```

Giving a power close too, but slight less than, a fixed-effects design.

- (e) Given these sample sizes, what is the smallest value of σ_τ^2/σ_e^2 that gives a (fixed-effects) 95% test a power of 0.90? (You will need to do this, and some the remaining problems, by trial and error.)

We try various values of σ_τ^2/σ_e^2 :

```
> 1-pf(2.578739, 5-1, 5*(10-1), 10*4*(1/2)) # trying 1/2
[1] 0.943019
1-pf(2.578739, 5-1, 5*(10-1), 10*4*(0.4)) # trying 0.4
[1] 0.8768408
1-pf(2.578739, 5-1, 5*(10-1), 10*4*(0.43)) # trying 0.43
```

[1] 0.9015374

Hence $\sigma_\tau^2/\sigma_e^2 = 0.43$, or (rearranging results from (a)),

$$\sigma_\tau^2/\sigma_T^2 = \frac{1}{1 + 1/(\sigma_\tau^2/\sigma_e^2)} = \frac{1}{1 + 1/0.43} = 0.30$$

Hence, this design has the power to detect an effect accounting for 30% or more of the total variation.

Note that we could also solve this more quickly by using the graphics commands in R, plotting power as a function of varying values of σ_τ^2/σ_e^2 . In particular to example power for $0.1 \leq \sigma_\tau^2/\sigma_e^2 \leq 0.9$, the R code is

```
> curve(1-pf(2.578739,5-1,5*(10-1),10*4*x), 0.1,0.9)
```

which returns a nice graph. We can focus in on finer regions of the curve by restricting the interval to smaller regions, e.g.,

```
> curve(1-pf(2.578739,5-1,5*(10-1),10*4*x), 0.4,0.5)
```

- (f) Given these sample sizes, what is the smallest value of σ_τ^2/σ_e^2 that gives a random-effects 95% test a power of 0.90?

Here, we need to solve for σ_τ^2/σ_e^2 in

$$\Pr \left[F_{N-1, N(n-1)} > \frac{2.578739}{1 + n(\sigma_\tau^2/\sigma_e^2)} \right] = 0.9$$

Using R [`qf(0.1, 5-1, 5*(10-1))`], we find that $\Pr [F_{N-1, N(n-1)} > 0.26323] = 0.9$, hence we solve for

$$\frac{2.578739}{1 + 10(\sigma_\tau^2/\sigma_e^2)} = 0.26323$$

giving $\sigma_\tau^2/\sigma_e^2 = 0.8796524$, so that σ_τ^2 must account for at least 46.8% of the total variance (using the expression in part (f) to covert σ_τ^2/σ_e^2 into σ_τ^2/σ_T^2 .)

- (g) How many replicates per factor are needed to give the fixed-effects ANOVA a power of 90% under a test of significant with $\alpha = 0.05$?

Once again, we can use trail and error, varying n to solve

$$\Pr(F_{5-1, 5(n-1), n(5-1)/9} \geq F_{5-1, 5(n-1), [0.95]}) = 0.9$$

in R, we first try $n = 30$,

```
> n <- 30; 1-pf(qf(0.95, 5-1, 5*(n-1)), 5-1, 5*(n-1), n*4/9)
```

```
[1] 0.8339192
```

```
n <- 35; 1-pf(qf(0.95, 5-1, 5*(n-1)), 5-1, 5*(n-1), n*4/9)
```

```
[1] 0.8940105
```

```
n <- 36; 1-pf(qf(0.95, 5-1, 5*(n-1)), 5-1, 5*(n-1), n*4/9)
```

```
[1] 0.9034603
```

$n = 36$ it is.

- (h) How many replicates per factor are needed to give the random-effects ANOVA a power of 90% under a test of significant with $\alpha = 0.05$? (Again, need to use trail and error)

Here we need to find n such that

$$\Pr \left[F_{5-1, 5(n-1)} > \frac{F_{5-1, 5(n-1), [0.95]}}{1 + n/9} \right] = 0.9$$

in R, we first try $n = 80$,

```
n <- 80; 1-pf(qf(0.95, 5-1, 5*(n-1))/(1+n/9), 5-1, 5*(n-1))
```

```

[1] 0.914335
> n <- 73; 1-pf(qf(0.95, 5-1,5*(n-1))/(1+n/9),5-1,5*(n-1))
[1] 0.9015613
> n <- 72; 1-pf(qf(0.95, 5-1,5*(n-1))/(1+n/9),5-1,5*(n-1))
[1] 0.8995078
giving n = 73.

```

2 : Optimal design for a random-effects ANOVA. Suppose you have a total $T = 100$ measurements that you can make, and you have to decide how best to allocate them over N and n in a random-effects design. Should one choose more factors (increase N) at the expense of fewer replicates n per factor? Obviously, there is some intermediate trade-off between the two. Suppose that the factor variance is $\sigma_\tau^2 = 10$ and the error variance $\sigma_e^2 = 20$.

(a) Compute the power of this design for the following combinations of N and n :

50,2 33,3 25,4 20,5 10,10 5,20 4,25 3,33 2,50

Hint: It might make sense to first write an R function to do this for arbitrary N, n

`fcrit <- function(N,n,alpha) qf(1-alpha,N-1,N*(n-1))` is the R code for the α -level critical value, $F_{N-1,N(n-1),[1-\alpha]}$ under the null hypothesis. Recall (Equation A5.30b in the power notes) that the power is

$$\Pr\left(F_{N-1,N(n-1)} > \frac{F_{N-1,N(n-1),[1-\alpha]}}{1 + n\sigma_\tau^2/\sigma_e^2}\right)$$

Using the above function, a new function, `fpower` can be written in R as

`fpower <- function(N,n,alpha,var) 1-pf(fcrit(N,n,alpha)/(1+n*var),N-1,N*(n-1))` ■

where `var = σ_τ^2/σ_e^2` . For example, `fpower(33,3,0.05,10/20)` returns `[1] 0.9107213`. Running through the above values of N, n ,

50,2 33,3 25,4 20,5 10,10 5,20 4,25 3,33 2,50
0.7805 0.9107 0.9472 0.9593 0.9626 0.9242 0.8962 0.8384 0.6980

(b) What is the optimal design (i.e., which combination of N and n gives the largest power)?

$N = n = 10$

(c) Repeat (a) and (b) assuming $\sigma_\tau^2 = 20, \sigma_e^2 = 10$

50,2 33,3 25,4 20,5 10,10 5,20 4,25 3,33 2,50
0.99995 0.99999 0.99999 0.99998 0.9996 0.9932 0.9838 0.9549 0.8459