

# Literature Cited

Numbers in brackets following each reference denote the Lectures in which the reference is cited.

- Akey, J. M., G. Zhang, K. Zhang, L. Jin, and M. D. Shriver. 2002. Interrogating a high-density SNP map for signatures of natural selection. *Genome Reserach* 12: 1805–1814. [8]
- Anisimova, M., J. P. Bielawski, and Z. Yang. 2001. Accuracy and power of the likelihood ratio test in detecting adaptive molecular evolution. *Mol. Biol. Evol.* 18: 1585–1592. [8]
- Arnold, S. J. 1986. Limits on stabilizing, disruptive, and correlational selection set by the opportunity for selection. *Amer. Nat.* 128: 143–146. [12, 14]
- Arnold, S. J., and M. J. Wade. 1984a. On the measurement of natural and sexual selection: theory. *Evolution* 38: 709–719. [12]
- Arnold, S. J., and M. J. Wade. 1984b. On the measurement of natural and sexual selection: applications. *Evolution* 38: 720–734. [12]
- Atkins, K. D. and R. Thompson. 1986. Predicted and realized response to selection for an index of bone length and body weight in Scottish Blackface sheep. 1. Responses in the index and component traits. *Anim. Prod.* 43: 421–435. [9]
- Beavis, W. D. 1994. The power and deceit of QTL experiments: lessons from comparative QTL studies. In *49th Annual Corn and Sorghum Research Conference*. pp. 252–268. American Seed Trade Association, Washington, D.C. [17]
- Beniwal, B. K., I. M. Hastings, R. Thompson, and W.G. Hill. 1992a. Estimation of changes in genetic parameters in selected lines of mice using REML with an animal model. 1. Lean mass. *Heredity* 69: 352–360. [10]
- Beniwal, B. K., I. M. Hastings, R. Thompson, and W.G. Hill. 1992b. Estimation of changes in genetic parameters in selected lines of mice using REML with an animal model. 2. Body weight, body composition and litter size. *Heredity* 69: 361–371. [10]
- Benjamini, Y. and D. Yekutieli. 2001. The control of the false discovery rate in multiple testing under dependency. *Annals of Statistics*. 29:1165-1188. [18]
- Blair, H. T., and E. J. Pollak. 1984. Estimation of genetic trend in a selected population with and without the use of a control population. *J. Animal Sci.* 58: 878–886. [10]
- Blows, M. W., and R. Brooks. 2003. Measuring nonlinear selection. *Amer. Natl.* 162: 815–820. [12, 14]
- Blows, M. W., S. F. Chenoweth, and E. Hine. 2004. Orientation of the genetic variance-covariance matrix and fitness surface for multiple male sexually selected traits. *Amer. Natl.* 163: 329–340. [13, 14]
- Boag, P. T., and P. R. Grant. 1981. Intense natural selection in a population of Dawrin’s finches (*Geospizinae*) in the Galápagos. *Science* 214: 82–84. [12]
- Booksten, F. L. 1989. Comment on a rate test. *Evolution* 43: 1569–1570. [7]
- Box, G. E. P. and N. R. Draper. 1987. *Empirical model-building and response surfaces*. John Wiley and Sons, New York. [14]
- Bradford, G. E., D. T. Torell, G. M. Spurlock, and W. C. Weir. 1963. Performance and variability of offspring of crossbred and purebred rams. *J. Animal Science* 22: 617–623. [6]
- Breden, F. J. and M. J. Wade. 1989. Selection within and between kin groups of the imported willow leaf beetle. *Amer. Nat.* 134: 35–50. [12]
- Brem, R. B., G. Yvert, R. Clinton, and L. Kruglyak. 2002. Genetic dissection of transcriptional regulation in budding yeast. *Science* 296: 752–755. [18]
- Brem, R., and L. Kruglyak. 2005. The landscape of genetic complexity across 5,700 gene expression traits in yeast. *Proc. Natl. Acad. Sci. USA* 102: 1572-1577. [18]
- Brodie, E. D. 1992. Correlational selection for color pattern and antipredator behavior in the garter snake *Thamnophis ordinoides*. *Evolution* 46: 1284–1298. [14]
- Brooks, R., and J. A. Endler. 2001. Direct and indirect sexual selection and quantitative genetics of male traits in guppies (*Poecilia reticulata*). *Evolution* 55: 1002 - 1015. Brooks and Endler (2001) [14]
- Brown, W. M. and A. E. Bell. 1961. Genetic analysis of plateaued populations of *Drosophila*. *Genetics* 46: 407–425. [11]
- Brown, W. M. and A. E. Bell. 1980. An experimental comparison of selection alternatives to plateaued response. *Genetics* 94: 477–496. [11]
- Bulmer, M. G. 1971. The effect of selection on genetic variability. *Amer. Nat.* 105: 201–211. [9]
- Bulmer, M. G. 1974. Linkage disequilibrium and genetic variability. *Genet. Res.* 23: 281–289. [9]
- Bulmer, M. G. 1976. The effect of selection on genetic variability: a simulation study. *Genet. Res.* 28: 101–117. [9]
- Bulmer, M. G. 1980. *The mathematical theory of quantitative genetics*. Oxford Univ. Press, NY. [9]
- Bumpus, H. 1899. The elimination of the unfit as illustrated by the introduced sparrow, *Passer domesticus*. Biol. Lectures, Marine Biol. Lab, Woods Hole: 209–226. [12]
- Burrows, P. M. 1972. Expected selection differentials for directional selection. *Biomet.* 28: 1091–1100. [9]
- Cantet, R. J. C. and C. Smith. 1991. Reduced animal model for marker assisted selection using best linear unbiased prediction. *Génét. Sél. Evol.* 23: 221–233. [13]
- Carmon, J. L., H. A. Stewart, C. C. Cockerham, and R. E. Comstock. 1956. Prediction equations for rotational crossbreeding. *J. Animal Sciences* 15: 930–936. [6]
- Cavalli-Sforza, L. L. 1966. Population structure and human evolution. *Prod. Royal Sco. London. Series B* 164: 362–379. [8]
- Charlesworth, B. 1984. The cost of phenotypic evolution. *Paleobiology* 10: 319–327. [15]
- Cheung, V. G., L. K. Conlin, T. M. Weber, M. Arcaro, J.-Y. Jen M. Morley, and R. S. Spielman. 2004. Natural variation in human gene expression asses in lymphoblastiod cells. *Nature Genetics* 33: 422- 42 [18]

## 2 LITERATURE CITED

- Cheverud, J. M. 1984. Quantitative genetics and developmental constraints on evolution by selection. *J. Theor. Biol.* 110: 155–171. [14]
- Clayton, G. A., and A. Robertson. 1957. An experimental check on quantitative genetic theory. II. The long term effects of selection. *J. Genet.* 55: 152–170. [11]
- Conant, G. C., A. Wagner. 2004. Duplicate genes and robustness to transient gene knock-downs in *Caenorhabditis elegans*. *Proc. R. Soc. Lond. B* 271: 89–96. [18]
- Connner, J. K. 2001. How strong is natural selection? *Trends in Ecology and Evolution* 16: 215–217. [12]
- Conover, W. J. 1980. *Practical nonparametric statistics* (2nd Edition). Wiley and Sons, New York. [12]
- Copeman, J. B., and 15 others. 1995. Linkage disequilibrium mapping of a type 1 diabetes susceptibility gene (IDDM7) to chromosomes 2q31-q33. *Nature Genetics* 9: 80–85. [16]
- Cotterman, C. W. 1940. A calculus for statistico-genetics. Ph. D. Thesis, Ohio State Univ., Columbus. [4]
- Crespi, B. J. and F. L. Bookstein. 1988. A path-analytic model for the measurement of selection on morphology. *Evolution* 43: 18–28. [14]
- Crow, J. F. 1958. Some possibilities for measuring selection intensities in man. *Hum. Biol.* 30: 1–13. [12]
- Crow, J. F., and M. Kimura. 1970. *An introduction to population genetics theory*. Harper and Row, Publ., New York. [11]
- Crow, J. F., and N. E. Morton. 1955. Measurement of gene frequency drift in small populations. *Evolution* 9: 202–214. [11]
- Cruz, A., and J. W. Wiley. 1989. The decline of an adaptation in the absences of a presumed selection pressure. *Evolution* 43: 55–62. [11]
- Cutter, A. D., and S. Ward. 2004. Sexual and temporal dynamics of molecular evolution in *C. elegans* development. *Mol. Biol. Evol.* 22: 178–188. [18]
- Damerval, C., A. Maurice, J. M. Josse, and D. de Vienne. 1994. Quantitative trait loci underlying gene product variation: a novel perspective by analyzing regulation of genome expression. *Genetics* 137: 289–301.
- Darwin, C. 1871. *The descent of man and selection in relation to sex*. John Murray, London. [12]
- Davies, J. L., and 15 others. 1994. A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 371: 130–136. [17]
- Dempster, E. R. 1955. Genetic models in relation to animal breeding. *Biometrics* 11: 535–536. [11]
- Denver, D. R., K. Morris, J. T. Streelman, S. K. Kim, M. Lynch, and W. K. Thomas. 2005. The transcriptional consequences of mutation and natural selection in *Caenorhabditis elegans*. *Nature Genetics* 37: 544–548 [18]
- Depaulis, F., and M. Veuille. 1998. Neutrality tests based on the distribution of haplotypes under an infinite-site model. *Mol. Biol. Evol.* 15: 1788–1790. [8]
- Devlin, B., and K. Roeder. 1999. Genomic control for association studies. *Biometrics* 55: 997–1004. [16]
- Devlin, B., and K. Roeder, and L. Wasserman. 2001. Genomic control, a new approach to genetic-based association studies. *Theor. Pop. Biol.* 60: 155–166. [16]
- Dickerson, G. E. 1955. Genetic slippage in response to selection for multiple objectives. *Cold Spring Harb Symp Quant Biol.* 20:213–24. [13]
- Dickerson, G. E. 1969. Techniques for research in quantitative animal genetics. In Am. Soc. Animal Sci., *Techniques and procedures in animal science research*, pp. 36–79. Albany, NY. [6]
- Dickerson, G. E., C. T. Blunn, A. B. Chapman, R. M. Kottman, J. L. Krider, E. J. Warwick, J. A. Whatley, Jr., M. L. Baker, J. L. Lush, and L. M. Winters. 1954. Evaluation of selection in developing inbred lines of swine. *Mo. Agric. Exp. Sta. Res. Bull.* 551. [13]
- Dobbs, K. S. 1955. Hybrid vigor in plant breeding. *Proced. Roy. Soc. London B* 144: 185–192. [6]
- Dowdeswell, W. H. 1961. Experimental studies on natural selection in the butterfly, *Maniola jurtina*. *Heredity* 16: 39–52. [12]
- Downhower, J. F., L. S. Blumer, and L. Brown. 1987. Opportunity for selection: an appropriate measure for evaluating variation in the potential for selection? *Evolution* 41: 1395–1400. [12]
- Dudley, J. W. 1977. 76 generations of selection for oil and protein percentage in maize. In E. Pollak, O. Kempthorne, and T. B. Bailey, Jr., (eds.), *Proceedings of the international conference on quantitative genetics*, pp. 459–473. Iowa State Univ. Press, Iowa. [11]
- Dudley, J. W., and R. J. Lambert. 1992. Ninety generations of selection for oil and protein in maize. *Maydica* 37: 81–87. [11]
- Duret, L. and D. Mouchiroud. 2000. Determinants of substitution rates in mammalian genes: expression pattern affects selection intensity but not mutation rate. *Mol. Biol. Evol.* 17: 68–74. [18]
- Duvick, D. N. 1999. Heterosis: feeding people and protecting natural resources. In J. G. Coors and S. Pandey (eds). *The genetics and exploitation of heterosis in crops*, pp. 19–29. American Society of Agronomy, Madison, Wisconsin. [6]
- Eisen, E. J. 1972. Long-term selection response for 12-day litter weight in mice. *Genetics* 72: 129–142. [11]
- Eisen, E. J. 1975. Population size and selection intensity effects on long-term selection response in mice. *Genetics* 79: 305–323. [11]
- Eisen, M. B., P.T. Spellman, P. O. Brown, and D. Botstein. 1998. Cluster analysis and display of genome-wide expression patterns. *PNAS* 95:14863–14868. [18]
- Endler, J. A., 1986. *Natural selection in the wild*. Princeton University Press, Princeton, New Jersey. [12]
- Endo, T., K. Ikeo, and T. Gojobori. 1996. Large-scale search for genes on which positive selection may operate. *Mol. Biol. Evol.* 13: 685–690. [8]
- Enfield, F. D. 1980. Long term effects of selection: The limits to response. In A. Roberston, (ed.), *Selection*

- experiments in laboratory and domestic animals, pp. 69–86. Commonwealth Agricultural Bureaux. [11]
- Ewens, W. J. 1972. The sampling theory of selectively neutral alleles. *Theor. Popul. Biol.* 3: 87–112. [8]
- Ewens, W. J., and R. S. Spielman. 2001. Locating genes by linkage and association. *Theor. Popul. Biol.* 60: 135–139. [16]
- Falconer, D. S. 1954. Asymmetrical response in selection experiments. In A. A. Buzzati-Traverso (ed.) Symposium on genetics of population structure pp. 16–41. International Union Biol. Sci., Naples Series B, No. 15. [10]
- Falconer, D. S. 1955. Patterns of response in selection experiments with mice. *Cold Spring Harbor Symp. Quant. Biol.* 20: 178–196. [11]
- Falconer, D. S. 1960. The genetics of litter size in mice. *J. Cell and Comp. Physiol.* 56 (Suppl 1): 153–167. [9]
- Falconer, D. S. 1965. Maternal effects and selection response. *Proc. XIth Internat. Cong. Genetics* 3: 763–774. [9]
- Falconer, D. S. 1971. Improvement of litter size in a strain of mice at a selection limit. *Genet. Res.* 17: 215–235. [11]
- Falconer, D. S. and J. W. B. King. 1953. A study of selection limits in the mouse. *J. Genet.* 51: 561–581. [11]
- Falconer, D. S. 1977. Some results of the Edinburgh selection experiments with mice. In E. Pollak, O. Kempthorne, and T. B. Bailey, Jr., (eds.), *Proceedings of the international conference on quantitative genetics*, pp. 101–115. Iowa State Univ. Press, Iowa. [11]
- Fay, J. C., H. L. McCullough, P. D. Sniegowski, and M. B. Eisen. 2004. Population genetic variation in gene expression is associated with phenotypic variation in *Saccharomyces cerevisiae*. *Genome Biology* 5(4): R26 [18]
- Fay, J. C., and C.-I. Wu. 2000. Hitchhiking under positive Darwinian selection. *Genetics* 155: 1405–1413. [8]
- Fell, D. 1997. *Understanding the Control of Metabolism*. Portland Press, London. [18]
- Felsenstein, J. 1977. Multivariate normal genetic models with a finite number of loci. In E. Pollak, O. Kempthorne, and T. B. Bailey, Jr., (eds.), *Proceedings of the international conference on quantitative genetics*, pp. 227–246. Iowa State Univ. Press, Iowa. [15]
- Felsenstein, J. 1983. Statistical inference of phylogenies (with discussion). *Journal Royal Statistical Society A*, 146:246–272. [18]
- Fernando, R. L. 1990. Statistical problems in marker-assisted selection for QTL. In, *Proceedings 4th World Congress Genetics Applied to Livestock Production* 13: 433–436. [13]
- Fernando, R. L. and M. Grossman. 1989. Marker-assisted selection using best linear unbiased prediction. *Genet. Sel. Evol.* 21: 467–477. [13]
- Fernando, R. L., D. Nettleton, B. Southey, J. C.M. Dekkers, M. F. Rothschild, and M. Soller 2004. Controlling the proportion of false positives (FP) in multiple dependent tests. *Genetics* 166: 611–619. [18]
- Fields, S., and O. Song. 1989. A novel genetic system to detect protein-protein interactions. *Nature* 340: 245–246. [18]
- Fisher, R. A. 1918. The correlation between relatives on the supposition of Mendelian inheritance. *Trans. Royal Soc. Edinburgh* 52: 399–433. [1,3,4]
- Fisher, R. A. 1922. On the dominance ratio. *Proc. Royal Society Edinburgh* 42: 321–341. [6]
- Fitch, W. M., R. M. Bush, C. A. Vender, and N. J. Cox. 1997. Long-term trends in the evolution of H(3)Ha1 human influenza type A. *Proc. Natl. Acad. Sci. USA* 94: 7712–7718. [8]
- Ford, M. J. 2002. Applications of selective neutrality tests to molecular ecology. *Mol. Ecology* 11: 1245–1262. [8]
- Frankham, R., B. H. Yoo and J. S. F. Barker. 1988. Reproductive fitness and artificial selection in animal breeding: culling on fitness prevents a decline in reproductive fitness in lines of *Drosophila melanogaster* selected for increased inebriation time. *Theor. Appl. Genet.* 76: 909–914. [11]
- Frankham, R., and R. K. Nurthen. 1981. Forging links between population and quantitative genetics. *Theor. Appl. Genet.* 59: 251–263. [11]
- Frahm, R. R. and K.-I. Kojima. 1966. Comparison of selection responses on body weight under divergent larval density conditions in *Drosophila pseudoobscura*. *Genetics* 54: 625–637. [11]
- Fredeen, H. T. 1984. Selection limits: have they been reached with pigs? *Can. J. Anim. Sci.* 64: 223–234. [11]
- Fu, Y.-X. 1996. New statistical tests of neutrality for DNA samples from a population. *Genetics* 143: 557–570. [8]
- Fu, Y.-X. 1997. Statistical tests of neutrality of neutrality against population growth, hitchhiking and background selection. *Genetics* 147: 915–925. [8]
- Fu, Y.-X., and W.-H. Li. 1993. Statistical tests of neutrality of mutations. *Genetics* 133: 693–709. [8]
- Gauch, H. G., Jr. 1992. *Statistical analysis of regional yield trials: AMMI analysis of factorial designs*. Elsevier, Amsterdam, the Netherlands. [18]
- Goddard, M. E. 1992. A mixed model for analysis of data on multiple genetic markers. *Theor. Appl. Genet.* 83: 878–886. [13]
- Goldman, N., and Z. Yang. 1994. A codon-based model of nucleotide substitution for protein-coding DNA sequences. *Mol. Biol. Evol.* 11: 725–736. [8]
- Gowe, R. S., A. Robertson, and B. D. H. latter. 1959. Environment and poultry breeding problems 5: The design of poultry control strains. *Poultry Sci* 38: 462–471. [6]
- Gowe, R. S. 1983. Lessons from selection studies in poultry for animal breeders. *Proc 32nd Annual Breed. Roundtable*, pp. 22–50. [11]
- Gowe, R. S., R. W. Fairfull, I. McMillan, and G. S. Schmidt. 1993. A strategy for maintaining high fertility and hatchability in a multiple-trait egg stock selection program. *J. Pul. Sci* 72: 1433–1448. [11]

#### 4 LITERATURE CITED

- Graham, A. 1981. *Kronecker products and matrix calculus with applications*. Halsted Press, New York. [15]
- Grant (1990-1) [7]**
- Gregory, K. E., J. C. M. Trail, H. J. S. Marples, and J. Kakonge. 1985. Heterosis and breed effects on maternal and individual traits of *Bos Indicus* breeds of cattle. *J. Animal Sci.* 60: 1175–1180. [6]
- Griffing, B. 1960a. Theoretical consequences of truncation selection based on the individual phenotype. *Aust. J. Biol. Sci.* 13: 307–343. [9]
- Griffing, B. 1960b. Accommodation of linkage in mass selection theory. *Aust. J. Biol. Sci.* 13: 501–526. [9]
- Griffiths, A. J. F., J. H. Miller, D. T. Suzuki, R. C. Lewontin, and W. M. Gelbart. 1996. *An introduction to genetic analysis*. 6th Ed. W. H. Freeman, NY. [12]
- Gu, Z., D. Nicolae, H. H. Lu, and W. H. Li. 2002. Rapid divergence in expression between duplicate genes inferred from microarray data. *Trend in Genetics* 18: 609–613. [18]
- Haldane, J. B. S. 1919. The combination of linkage values, and the calculation of distance between the loci of linked factors. *J. Genetics* 8: 299–309. [3, 16]
- Haldane, J. B. S. 1954. The measurement of natural selection. *Proc. IX Internat. Cong. Genet.* 1: 480–487. [15]
- Hanrahan, J. P., E. J. Eisen and J. E. Legates. 1973. Effects of population size and selection intensity on short-term response to selection for postweaning gain in mice. *Genetics* 73: 513–530. [11]
- Harter, H. L. 1961. Expected values of normal order statistics. *Biometrika* 48: 151–166. [9]
- Hartley, H. O., and J. N. K. Rao. 1967. Maximum-likelihood estimation for the mixed analysis of variance model. *Biometrika* 54: 93–108. [5]
- Harvey, W. R. and G. D. Bearden. 1962. Tables of expected genetic progress in each of two selected traits. USDA Ag. Res. Service Publication no 20-12. [13]
- Haseman, J. K., and R. C. Elston. 1972. The investigation of linkage between a quantitative trait and a marker locus. *Behav. Genet.* 2: 3–19. [17]
- Hästbacka, J., A. de la Chapelle, I. Kaitila, P. Sistonen, A. Weaver, and E. Lander. 1992. Linkage disequilibrium mapping in isolated founder populations: diastrophic dysplasia in Finland. *Nature Genetics* 2: 204–211. [16]
- Hazel, L. N. 1943. The genetic basis for constructing selection indexes. *Genetics* 28: 476–490. [13]
- Heath, S. C., G. Bulfield, R. Thompson, and P. D. Keightley. 1995. Rates of change of genetic parameters of body weight in selected mouse lines. *Genet. Res. Camb.* 66: 19–25. [10]
- Henderson, C. R. 1950. Estimation of genetic parameters. *Ann. Math. Stat.* 21: 309–310. [5]
- Henderson, C. R. 1953. Estimation of variance and covariance components. *Biometrics* 9: 226–252. [5]
- Henderson, C. R. 1963. Selection index and the expected genetic advance In W. D. Hanson and H. F. Robinson (eds.), *Statistical genetics and plant breeding*, pp. 141–163. Natl. Acad. Sci., Natl. Res. Council Publ. No. 982, Washington, D. C. [5]
- Henderson, C. R. 1973. Sire evaluation and genetic trends. In Proceedings of the animal breeding and genetics symposium in honor of Dr. J. L. Lush, pp. 10–41. American Society of Animal Science, Champaign, IL. [5]
- Henderson, C. R. 1984. *Applications of linear models in animal breeding*. Univ. Guelph, Guelph, Ontario. [5]
- Hill, W. G. 1976. Order statistics of correlated variables and implications in genetic selection programmes. *Biometrics* 32: 889–902. [9]
- Hill, W. G. 1977. Order statistics of correlated variables and implications in genetic selection programmes. II. Response to selection *Biometrics* 33: 703–712. [9]
- Hill, W. G. 1982a. Rates of change in quantitative traits from fixation of new mutations. *Proc. Natl. Acad. Sci.* 79: 142–145. [11]
- Hill, W. G. 1982b. Predictions of response to artificial selection from new mutations. *Genet. Res.* 40: 255–278. [11]
- Hoekstra H., J. Hoekstra, D. Berrigan, S. Vignieri, C. Hill, A. Hoang, P. Gibert, P. Beerli, and J. G. King-solver. 2001. The strength and tempo of directional selection in the wild. *PNAS* 98: 9157–9160. [12]
- Hofmeyr, J.-H. S. and H. V. Westerhoff. 2001. Building the cellular puzzle. Control in multi-level reaction networks. *J. Theoret. Biol.* 208:261–285. [18]
- Hopkins, C. G. 1899. Improvement in the chemical composition of the corn kernel. *Ill. Agric. Exp. Stn. Bull.* 55: 205–240. [11]
- Howard, R. D. 1979. Estimating reproductive success in natural populations. *Amer. Nat.* 114: 221–231. [12]
- Hudson, R. R., M. Kreitman, and M. Aguade. 1987. A test of neutral molecular evolution based on nucleotide data. *Genetics* 116: 153–159. [8]
- Hughes, A. L., and M. Nei. 1998. Pattern of nucleotide substitution at major histocompatibility complex class I loci reveals overdominant selection. *Nature* 335: 167–170. [8]
- Hunton, P. 1984. Selection limits: have they been reached in the poultry industry? *Can. J. Anim. Sci.* 64: 217–221. [11]
- James, J. W. 1962. Conflict between directional and centripetal selection. *Heredity* 17: 487–499. [11]
- James, J. W. 1965. Response curves in selection experiments. *Heredity* 20: 57–63. [11]
- Janssen, G. M., D. DeJong, E. N. G. Joosse, and W. Scharloo. 1988. A negative maternal effect in springtails. *Evolution* 42: 828–834. [9]
- Jenkin, F. 1867. Origins of species. *North British Review* 46: 277–318. [3]
- Jeong, H., B. Tombor, R. Albert, Z. N. Oltvai, and A.-L. Barabasi. 2000. The large-scale organization of metabolic networks. *Nature* 407: 651–654. [18]
- Jones, D. F. 1918. The effects of inbreeding and crossbreeding upon development. *Conn. Agric. Exp. Sta. Bull.* 207: 5–100. [6]
- Jones, D. F. 1922. The production of single and double first-generation corn hybrids. *J. American Agrn.*

- Society* 14: 241–252. [6]
- Jones, D. F., and H. L. Everett. Hybrid field corn. *Bull.* 532. Conn. Agric. Exp. Stn. [6]
- Jones, L. P., R. Frankham and J. S. F. Barker. 1968. The effects of population size and selection intensity in selection for a quantitative character in *Drosophila*. II. Long-term response to selection. *Genet. Res.* 12: 249–266. [11]
- Jordan, I. K., L. Marino-Ramirez, and E. V. Koonin. 2005. Evolutionary significance of gene expression divergence. *Gene*, 345: 19–126. [18]
- Jordan, N. 1991. Multivariate analysis of selection in experimental populations derived from hybridization of two ecotypes of the annual plant *Diodia teres*. *Evolution* 45: 1760–1772. [14]
- Kacser, H., and J. A. Burns. 1973. The control of flux. *Symp. Soc. Experimental Biology* 27: 65–104. [18]
- Kacser, H. and J. A. Burns. 1981. The molecular basis of dominance. *Genetics* 97:639–666. [18]
- Kalisz, S. 1986. Variable selection on the timing of germination in *Collinsia verna* (Scrophulariaceae). *Evolution* 40: 479–491. [14]
- Kaplan, N., and R. Morris. 2001. Prospects for association-based fine mapping of a susceptibility gene for a complex disease. *Theor. Pop. Biol.* 60: 18–191. [16]
- Karlin, S., and H. M. Taylor. 1981. *A second course in stochastic processes*. Academic Press, New York. [7]
- Karn, M. N., and L. S. Penrose. 1951. Birth weight and gestation time in relation to maternal age, parity and infant survival. *Ann. Eugen.* 16: 147–164. [12]
- Kauffman, S.A. 1969. Homeostasis and Differentiation in Random Genetic Control Networks. *Nature*. 224: 177–178. [18]
- Kayser, M., S. Brauer, and M. Stoneking. 2003. A genome scan to detect candidate regions influenced by local natural selection in human populations. *Mol. Biol. Evol.* 20: 893–900. [8]
- Kendall, M., and A. Stuart. 1983. *The advanced theory of statistics. Volume 3: Design and analysis, and time-series*. 4th Ed. Griffin, London. [14]
- Kennedy, B. W. 1984. Selection limits: have they been reached with the dairy cow? *Can. J. Anim. Sci.* 64: 207–215. [11]
- Kerr, M.K. and G.A. Churchill. 2001. Analysis of variance for gene expression microarray data. *J. Comp. Biol.* 7:819–837. [18]
- Khaitovich, P., G. Weiss, M. Lachmann, I. Hellmann, W. Enards, B. Muetzel, U. Wirkner, W. Ansorge, and S. Pääbo. 2004. A neutral model of transcriptome evolution. *PLoS Biology* 2: 682–689. [18]
- Khaitovich, P., I. Hellmann, W. Enards, K. Nowick, M. Leinweber, H. Franz, G. Weiss, M. Lachmann, and S. Pääbo. 2005. Parallel patterns of evolution in the genomes and transcriptomes of Humans and Chimpanzees. 2005. *Science* 309: 1850–1854. [18]
- Kidder, R. W., M. Koger, J. H. Meade, and J. R. Crockett. 1964. Systems of crossbreeding for beef production in Florida. *Bull. Agric. Experimental Stations, Florida University* No. 673, 19 pp. [6]
- Kim, Y., and W. Stephan. 2002. Detecting a local signature of genetic hitchhiking along a recombining chromosome. *Genetics* 160: 765–777. [8]
- Kimura, M. 1957. Some problems of stochastic processes in genetics. *Ann. Math. Stati.* 28: 882–901. [3]
- Kimura, M. 1968. Evolutionary rate at the molecular level. *Nature* 217: 624–626. [3]
- Kimura, M. 1983. *The neutral theory of molecular evolution*. Cambridge Univ. Press, UK. [7, 8]
- Kimura, M., and J. F. Crow. 1964. The number of alleles that can be maintained in a finite population. *Genetics* 49: 725–738. [3,7]
- Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. E. Hill, A. Hoang, P. Gilbeert, and P. Beerli. 2001. The strength of phenotypic selection in natural populations. *Amer. Natl.* 157: 245–261. [12, 14]
- Knight, G. R. and A. Robertson. 1957. Fitness as a measurable character in *Drosophila*. *Genetics* 42: 524–530. [11]
- Knowler, W. C., R. C. Williams, D. J. Pettitt, and A. G. Steinberg. 1988. *Gm<sup>3,5,13,14</sup>* and type 2 diabetes mellitus: an association in American Indians with genetic admixture. *Am. J. Hum. Genet.* 43: 520–526. [14]
- Kosambi, D. D. 1944. The estimation of map distances from recombination values. *Ann. Eugen.* 12: 172–175. [3, 16]
- Kownacki, M. 1979. Effect of reciprocal crossing of selected lines of mice. *Theor. Appl. Genet.* 54: 169–175. [11]
- Krzanowski, W. J. 1979. Between-group comparisons of principal components. *J. Amer. Stat. Assoc.* 74: 703–707. [14]
- Kruuk, L. E. B., J. Slate, J. M. Pemberton, S. Brotherstone, F. Guinness, T. Clutton-Brock. 2002. Antler size in red deer: Heritability and selection but no evolution. *Evolution* 56: 1683–1695. [15]
- Lande, R. 1976. Natural selection and random genetic drift in phenotypic evolution. *Evolution* 30: 314–334. [7, 12, 14, 15]
- Lande, R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry. *Evolution* 33: 402–416. [13, 14, 15]
- Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters. *Evolution* 37: 1210–1226. [12, 14]
- Lande, R., and R. Thompson. 1990. Efficiency of marker-assisted selection in the improvement of quantitative traits. *Genetics* 124: 743–756. [13]
- Lander, E. S., and D. Botstein. 1989. Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* 121: 185–199 (Correction 136: 705). [17]
- Latter, B. D. H. 1960. Natural selection for an intermediate optimum. *Aust. J. Biol. Sci.* 13: 30–35. [11]
- Latter, B. D. H. 1965. The response to artificial selection due to autosomal genes of large effect. I. Changes

- in gene frequency at an additive locus. *Aust. J. Biol. Sci.* 18: 585–598. [11]
- Lee, K.E., N. Sha, E.R. Dougherty, M. Vannucci, and B.K. Mallick. 2003. Gene selection: a Bayesian variable selection approach. *Bioinformatics* 19:90–97. [18]
- Lemos, B., C. D. Meiklejohn, M. Caceres, and D. L. Hartl. 2005. Rates of divergence in gene expression profiles of primates, mice, and flies: stabilizing selection and variability among functional categories. *Evolution* 59: 126–137. [18]
- Lerner, I. M. 1950. *Population genetics and animal improvement*. Cambridge. [11]
- Lerner, I. M. 1954. *Genetic homeostasis*. Oliver and Boyd, London. [11]
- Lerner, I. M. and E. R. Dempster. 1951. Attenuation of genetic progress under continued selection in poultry. *Heredity* 5: 75–94. [11]
- Lewontin, R. C., and J. Krakauer. 1973. Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. *Genetics* 74: 175–195. [8]
- Liao, J.C. and J. Delgado. 1997. Flux calculation using control constraints. *Biotechnology Progress* 14: 554–560. [18]
- Lin, C. Y., and F. R. Allaire. 1977. Heritability of a linear combination of traits. *Theor. Appl. Genet.* 51: 1–3. [13]
- Lynch, M. and B. Walsh. 1998. *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Sunderland, MA. [7]
- Mackay, T. F. C. 1985. Transposable element-induced response to artificial selection in *Drosophila melanogaster*. *Genetics* 111: 351–374. [10]
- Maddox, G. D. and J. Antonovics. 1983. Experimental ecological genetics in *Plantago*: a structural equation approach to fitness components in *P. aristata* and *P. patagonica*. *Ecology* 64: 1092–1099. [14]
- Magee and Hazel (1959) [6]**
- Makova, K. D. and W. H. Li. 2003. Divergence in the spatial pattern of gene expression between human duplicate genes. *Genome Res.* 14: 1610–1616. [18]
- McDonald, J. H., and M. Kreitman. 1991. Adaptive protein evolution at the *Adh* locus in *Drosophila*. *Nature* 351: 652–654. [8] [8]
- McDowell, R. E. 1981. Limitations for dairy production in developing countries. *J. Dairy Sci.* 64:2463–2475. [6]
- McDowell, R. E. 1985. Crossbreeding in tropical areas with emphasis on milk, health, and fitness. *J. Dairy Sci.* 68: 2418–2435. [5]
- McGuigan, K., S. F. Chenoweth, and M. W. Blows. 2005. Phenotypic divergence along lines of genetic variance. *Amer. Nat.* 165: 32–43. [13]
- Meyer, K., and W. G. Hill. 1991. Mixed model analysis of a selection experiment for food intake in mice. *Genet. Res. Camb.* 57: 71–81. [10]
- Mitchell-Olds, T. 1987. Analysis of local variation in plant size. *Ecology* 68: 82–87. [14]
- Mitchell-Olds, T., and J. Bergelson. 1990. Statistical genetics of *Impatiens capensis*. II. Natural selection. *Genetics* 124: 417–421. [14]
- Mitchell-Olds, T., and R. G. Shaw. 1987. Regression analysis of natural selection: Statistical inference and biological interpretation. *Evolution* 41: 1149–1161. [12, 14]
- Morley, M., C. M. Molony, T. M. Weber, J. L. Devlin, K. G. Ewens, R. S. Ewens, R. S. Spielman, and V. G. Cheung. 2004. Genetic analysis of genome-wide variation in human gene expression. *Nature* 430: 743–747. [18]
- Morrison, D. F. 1976. *Multivariate statistical methods*. McGraw-Hill, New York. [15]
- Morton, N. E. 1955. Sequential tests for the detection of linkage. *Am. J. Hum. Genet.* 7: 277–318. [3, 17]
- Muse, S. V., and B. S. Gaut. 1999. A likelihood approach for comparing synonymous and nonsynonymous nucleotide substitution rates, with application to the chloroplast genome. *Mol. Biol. Evol.* 11: 715–724. [8]
- Newman, J. A., G. W. Rahnefeld and H. T. Fredeen. 1973. Selection intensity and response to selection for yearling weight in beef cattle. *Can. J. Anim. Sci.* 53: 1–12. [10]
- Nicholas, F. W. 1980. Size of population required for artificial selection. *Genet. Res.* 35: 85–105. [10]
- Nicholas, F. W. and A. Robertson. 1980. The conflict between natural and artificial selection in finite populations. *Theor. Appl. Genet.* 56: 57–64. [11]
- Nielsen, R. 2001. Statistical tests of selective neutrality in the age of genomics. *Heredity* 86: 641–647. [8]
- Nielsen, R. 2005. Molecular signatures of natural selection. *Annual Review of Genetics* 39: 197–218. [8]
- Nielsen, R., and Z. Yang. 1998. Likelihood models for detecting positively selected amino acid sites and applications to the HIV-1 envelope gene. *Genetics* 148: 929–936. [8]
- Neimann-Sørensen, A. and A. Robertson. 1961. The association between blood groups and several production characteristics in three Danish cattle breeds. *Acta Agric. Scand.* 11: 163–196. [13]
- Nitter, G. 1978. Breeding utilisation for meat production in sheep. *Animal Breeding Abstracts* 46: 131–143. [6]
- Nordskog, A. W. 1978. Some statistical properties of an index of multiple traits. *Theor. Appl. Genet.* 52: 91–94. [13]
- O'Donald, P. 1968. Measuring the intensity of selection. *Nature* 220: 197–198. [12]
- O'Donald, P. 1970. Change of fitness by selection for a quantitative character. *Theor. Pop. Biol.* 1: 219–232. [12]
- O'Donald, P. 1971. Natural selection for quantitative characters. *Heredity* 27: 137–153. [12]
- Pal, C., B. Papp, and L. D. Hurst. 2001. Highly expressed genes in yeast evolve slowly. *Genetics* 158: 927–931. [18]
- Parmigiani, G., E. S. Garrett, R. A. Irizarry, and S. L. Zeger (Eds). 2003. *The Analysis of Gene Expression Data*. Springer, New York. [18]

- Pearson, K. 1903. Mathematical contributions to the theory of evolution. XI. On the influence of natural selection on the variability and correlation of organs. *Phil. Trans. Roy. Soc. Lond. A* 200: 1–66. [12]
- Phillips, P. C. and S. J. Arnold. 1989. Visualizing multivariate selection. *Evolution* 43: 1209–1222. [12, 14]
- Pomp, D., M. Allan, and S. Wesolowski. 2004. Quantitative Genomics: Exploring the genetic architecture of complex trait predisposition. *J. Anim. Sci* 82: E300–E312. [18]
- Price, G. R. 1970. Selection and covariance. *Nature* 227: 520–521. [12]
- Price, G. R. 1972. Extension of covariance selection mathematics. *Ann. Hum. Genet* 35: 485–490. [12]
- Price, T. D., P. R. Grant, H. L. Gibbs, and P. T. Boag. 1984. Recurrent patterns of natural selection in a population of Darwin's finches. *Nature* 309: 787–789. [12]
- Price, T. D., and P. R. Grant. 1984. Life history traits and natural selection for small body size in a population of Darwin's finches. *Evolution* 38: 483–494. [12]
- Pritchard, J. K., and N. A. Rosenberg. 1999. Use of unlinked genetic markers to detect population stratification in association studies. *Amer. J. Hum. Genet* 65: 220–228. [16]
- Prout, T. 1965. The estimation of fitness from genotypic frequencies. *Evolution* 19: 546–551. [12]
- Prout, T. 1969. The estimation of fitness from population data. *Genetics* 63: 949–967. [12]
- Rausher, M. D. 1992. The measurement of selection on quantitative traits: biases due to environmental covariances between traits and fitness. *Evolution* 46: 616–626. [12, 14]
- Rawlings, J. O. 1976. Order Statistics for a special class of unequally correlated multinormal variates. *Biometrics* 32: 875–887. [9]
- Reeve, E. C. R. and F. W. Robertson. 1953. Studies in quantitative inheritance. II. Analysis of a strain of *Drosophila melanogaster* selected for long wings. *J. Genet.* 51: 276–316. [11]
- Rendel, J. M. 1943. Variations in the weights of hatched and unhatched duck's eggs. *Biometrika* 33: 48–58. [9]
- Reyment, R. A. 1982. Phenotypic evolution in a cretaceous foraminifer. *Evolution* 36: 1182–1199. [7]
- Richey, F. D. 1922. The experimental basis for the present status of corn breeding. *J. Am. Soc. Agron.* 14: 1–17. [6]
- Rifkin, S. A., J. Kim, and K. P. White. 2003. Evolution of gene expression in the *Drosophila melanogaster* subgroup. *Nature Genetics* 33: 138–144. [18]
- Risch, N. 1990a. Linkage strategies for genetically complex traits. II. The power of affected relative pairs. *Am. J. Hum. Genet.* 46: 229–241. [17]
- Risch, N. 1990b. Linkage strategies for genetically complex traits. III. The effect of marker polymorphism on analysis of affected relative pairs. *Am. J. Hum. Genet.* 46: 242–253 (Correction 51: 673–675). [17]
- Riska, B., T. Prout, and M. Turelli. 1989. Laboratory estimates of heritabilities and genetic correlations in nature. *Genetics* 123: 865–871. [5]
- Roberts, R. C. 1966. The limits to artificial selection for body weight in the mouse. II. The genetic nature of the limits. *Genet. Res.* 8: 361–375. [11]
- Robertson, A. 1956. The effect of selection against extreme deviants based on deviation or homozygosis. *J. Genet.* 54: 236–248. [11]
- Robertson, A. 1960. A theory of limits in artificial selection. *Proc. Royal Soc. B* 153: 234–249. [11]
- Robertson, A. 1966. A mathematical model of the culling process in dairy cattle. *Animal Production* 8: 95–108. [12]
- Robertson, A. 1977. Artificial selection with a large number of linked loci. In E. Pollak, O. Kempthorne, and T. B. Bailey, Jr. (eds.), *Proceedings of the international conference on quantitative genetics*, pp. 307–322. Iowa State Univ. Press, Ames. [11]
- Robertson, F. W. 1955. Selection response and the properties of genetic variation. *Cold Spring Harbor Symp. Quant. Biol.* 20: 166–177. [11]
- Rogers, J. S., and J. R. Edwardson. 1952. The utilization of cytoplasmic male-sterile hybrids in the production of corn hybrids. *Agronomy Journal* 44: 8–13. [6]
- Saxton, A.M. 1988. Further approximations for selection intensity. *Theor. Appl. Genet.* 76: 465–466. [9]
- Schadt, E. E., S. A. Monks, T. A. Drake, A. J. Lusis, N. Che, V. Collnayo, T. G. Ruff, S. B. Milligan, J. R. Lamb, G. Cavet, P. S. Linsley, M. Mao, R. B. Stoughton, and S. H. Friend. 2003. Genetics of gene expression surveyed in maize, mouse and man. *Nature* 422: 297–302. [18]
- Scheiner, S. M. 1989. Variable selection along a successional gradient. *Evolution* 43: 548–562. [14]
- Schluter, D. 1988. Estimating the form of natural selection on a quantitative trait. *Evolution* 42: 849–861. [12]
- Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. *Evolution* 50: 1766–1774. [13]
- Schluter, D., and D. Nychka. 1994. Exploring fitness surfaces. *Amer. Natl.* 143: 597–616. [12]
- Searle, S. S. 1982. *Matrix algebra useful for statistics*. John Wiley and Sons, New York. [15]
- Searle, S. R., G. Casella, and C. E. McCulloch. 1992. *Variance components*. John Wiley and Sons, NY. [5]
- Sheridan, A. K. 1988. Agreement between estimated and realised genetic parameters. *Animal Breeding Abstracts* 56: 877–889. [10]
- Shull, G. H. 1908. The composition of a field of maize. *Rpt. Am. Breed. Assoc.* 4: 296–301. [6]
- Shull, G. H. 1909. A pure-line method in corn breeding. *American Breeders' Association* 5: 51–59. [6]
- Simms, E. L. 1990. Examining selection on the multivariate phenotype: plant resistance to herbivores. *Evolution* 44: 1177–1188. [14]
- Simms, E. L. and M. D. Rausher. 1993. Patterns of selection on phytophage resistance in *Ipomoea purpurea*. *Evolution* 47: 970–976. [14]
- Simonsen, K. L., G. A. Churchill, and C. F. Aquadro. 1995. Properties of statistical tests of neutrality for DNA polymorphism data. *Genetics* 141: 413–429. [8]

- Small, J. R., and H. Kacser. 1993. Responses of metabolic systems to large changes in enzyme activities and effectors. 1. The linear treatment of unbranched Chains. *Eur. J. Biochemistry* 213: 613–624. [18]
- Smith, C. 1967. Improvement of metric traits through specific genetic loci. *Anim. Prod.* 9: 349–358. [13]
- Smith, C. 1969. Optimum selection procedures in animal breeding. *Anim. Prod.* 11: 433–442. [9]
- Smith, H. F. 1936. A discriminant function for plant selection. *Ann. Eugen.* 7: 240–250. [13]
- Smith, L. H. 1908. Ten generations of corn breeding. *Ill. Agric. Exp. Stn. Bull.* 128: 457–575. [11]
- Sorensen, D. A. and W. G. Hill 1982. Effect of short term directional selection on genetic variability: experiments with *Drosophila melanogaster*. *Heredity* 48: 27–33. [9]
- Sorensen, D. A., and B. W. Kennedy. 1984. Estimation of response to selection using least-squares and mixed model methodology. *J. Anim. Sci.* 58: 1097–1106 [10]
- Spielman, R. S., R. E. McGinniis, and W. J. Ewens. 1993. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am. J. Hum. Genet.* 52: 506–516. [16]
- Sprague, G. F., and L. A. Tatum. 1942. General vs. specific combining ability in single crosses of corn. *J. Am. Soc. Agron.* 34: 923–932. [6]
- Stewart, S. C. and D. J. Schoen. 1987. Patterns of phenotypic variability and fecundity selection in a natural population of *Impatiens pallida*. *Evolution* 41: 1290–1301. [14]
- Stratton, D. A. 1992. Life-cycle components of selection in *Erigeron annuus*: I. Phenotypic selection. *Evolution* 46: 92–106. [14]
- Storey, J. D., and R. Tibshirani. 2003a. SAM thresholding and false discovery rates for detecting differential gene expression in DNA microarrays. In G. Parmigiani, E. S. Garrett, R. A. Irizarry, and S. L. Zeger (Eds), *The Analysis of Gene Expression Data*, pp. 272–290. Springer, New York. [18]
- Storey, J. D., and R. Tibshirani. 2003b. Statistical significance for genomewide studies. *PNAS* 100: 9440–9445. [18]
- Storz, J. G., B. A. Payseur, and M. W. Nachman. 2004. Genomic scans of DNA variability in humans reveal evidence for selective sweeps outside of Africa. *J. Mol. Biol. Evol* 21: 1800–1811. [8]
- Subramanian, S., and S. Kumar. 2004. Gene expression intensity shapes evolutionary rates of the proteins encoded by the vertebrate genome. *Genetics* 168: 373–381. [18]
- Suzuki, Y., and T. Gojobori. 1999. A method for detecting positive selection at single amino acid sites. *Mol. Biol. Evol* 16: 1315–1328. [8]
- Suzuki, Y., and M. Nei. 2002. Simulation study of the reliability and robustness of the statistical methods for detecting positive selection at single amino acid sites. *Mol. Biol. Evol* 19: 1865–1869. [8]
- Sved, J. A. 1989. The measurement of fitness in *Drosophila*. In W. G. Hill, and T. F. C. Mackay, (eds.), *Evolution and animal breeding*, pp. 113–120. C. A. B. International, Wallingford. [12]
- Tajima, F. 1989. Statistical methods for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123: 585–595. [8]
- Tallis, G. M. 1987. Ancestral covariance and the Bulmer effect. *Theor. Appl. Genet.* 73: 815–820. [15]
- Tallis, G. M. and P. Leppard. 1988. The joint effects of selection and assortative mating on multiple polygenic characters. *Theor. Appl. Genet.* 75: 278–281. [15]
- Tamayo, P., D. Slonim, J. Mesirov, Q. Zhu, S. Kitareewan, E. Dmitrovsky, E. S. Lander, and T. R. Golub. 1999. Interpreting patterns of gene expression with self-organizing maps: Methods and application to hematopoietic differentiation. *PNAS* 96:2907–2912. [18]
- Tatum, L. A. 1971. The Southern Corn Leaf Blight epidemic. *Science* 171: 1113–1116. [6]
- Tavazoie, S., D. Hughes, M. J. Campbell, R. J. Cho, and G. M. Church. 1999. Systematic determination of genetic network architecture. *Nature Genetics* 22: 2816–285. [18]
- Thoday, J. M. and T. B. Boam. 1961. Regular responses to selection. I. Description of response. *Genet. Res.* 2: 161–176. [11]
- Thoday, J. M., J. B. Gibson, and S. G. Spickett. 1964. Regular responses to selection. II. Recombination and accelerated response. *Genet. Res.* 5: 1–19. [11]
- Thompson, R. 1986. Estimation of realized heritability in a selected population using mixed model methods. *Genet. Sel. Evol.* 18: 475–484. [10]
- Trail, J. C. M., K. E. Gregory, H. J. S. Marples, and J. Kakong. 1985. Comparison of Bos Tarus-Bos Indicus breed crosses with straightbred Bos Indicus breeds of cattle for maternal and individual traits. *J. Animal Sci.* 60: 1181–1187. [6]
- Trail, P. W. 1985. The intensity of selection: intersexual and interspecific comparisons require consistent measures. *Amer. Nat.* 126: 434–439. [12]
- Trustring, G. B. 1961. The correlations between relatives in a random mating diploid population. *Proc. Cambridge Phil. Soc.* 57: 315–320. [4]
- Ullstrup, A. J. 1972. The impacts of the southern corn leaf blight epidemics of 1970–71. *Annual Review of Phytopathology* 10: 37–50. [6]
- van der Werf, J. H. J., and I. J. M. de Boer. 1990. Estimation of additive genetic variance when base populations are selected. *J. Anim. Sci.* 68: 3124–3132. [10]
- Victoria, J. K. 1972. Clutch characteristics and egg discrimination ability of the African Village Weaver *Ploceus cucullatus Ibis* 114: 367–376. [11]
- Wade, M. J. and S. Kalisz. 1990. The causes of natural selection. *Evolution* 44: 1947–1955. [12]
- Wagner, A. Decoupled evolution of coding regions and mRNA expression patterns after gene duplication: implications for the neutralist-selectionist debate. *Proc.*

- Natl. Acad. Sci. USA* 97: 6579–6584. [18]
- Wall, J. D., and R. R. Hudson. 2001. Coalescent simulations and statistical tests of neutrality. *Mol. Biol. Evol.* 18: 1134–1135. [8]
- Wang, J. 1997. More efficient breeding systems for controlling inbreeding and effective population size in animals. *Heredity* 79: 591–599. [6]
- Wang, E. T., G. Kodama, P. Baldi, and R. K. Moyzis. 2006. Global landscape of recent inferred Darwinian selection for *Homo sapiens*. *Proc. Natl. Acad. Sci. USA* 103: 135–140. [8]
- Waterson, G. A. 1975. On the number of segregation sites. *Theor. Popul. Biol.* 7: 256–276. [8]
- Weber, K. E. 1990. Increased selection response in larger populations. I. Selection for wing-tip height in *Drosophila melanogaster* at three population sizes. *Genetics* 125: 579–584. [11]
- Weber, K. E. 1996. Large genetic changes at small fitness cost in large populations of *Drosophila melanogaster* selected for wind tunnel flight: rethinking fitness surfaces. *Genetics* 144: 205–213. [11]
- Weber, K. E. and L. T. Diggins. 1990. Increased selection response in larger populations. II. Selection for ethanol vapor resistance in *Drosophila melanogaster* at two population sizes. *Genetics* 125: 585–597. [11]
- Weldon, W. F. R. 1895. Attempt to measure the death-rate due to the selective destruction of *Carcinus moenas* with respect to a partial dimension. *Proc. Roy. Soc.* 57: 360–379. [15]
- Weller, J. I., Y. Kashi, and M. Soller. 1990. Power of daughter and granddaughter designs for determining linkage between marker loci and quantitative trait loci in dairy cattle. *J. Dairy Sci.* 73: 2525–2537. [17]
- Whitlock, M. C., and K. Fowler. 1999. The changes in genetic and environmental variance with inbreeding in *Drosophila melanogaster*. *Genetics* 152: 345–353. [6]
- Willis, J. H. 1993. The measurement of selection in partially inbred populations: An example from *Mimulus guttatus*. *Evolution* Submitted MS 666. [12, 14]
- Wilson, S. P., H. D. Goodale, W. H. Kyle, and E. F. Godfrey. 1971. Long term selection for body weight in mice. *J. Heredity* 62: 228–234. [11]
- Wong, W. S. W., Z. Yang, N. Goldman, and R. Nielsen. 2004. Accuracy and power of statistical methods for detecting adaptive evolution in the protein coding sequencess and for identifying positively selection sites. *Genetics* 168: 1041–1051. [8]
- Wright, S. 1922. Coefficients of inbreeding and relationship. *Am. Nat.* 56: 330–339. [4, 6]
- Wright, S. 1931. Evolution in mendelian populations. *Genetics* 16: 97–159. [7]
- Wright, S. 1938. Size of population and breeding structure in relation to evolution. *Science* 87: 430–431. [7]
- Wright, S. 1939. Statistical genetics in relation to evolution. *Actualités Scientifiques et Industrielles, No. 802. Exposé de Biometrie et de Statistique Biologique. XIII.* Hermann and Cie, Paris. [7]
- Wright, S. I., C. B. K. Yau, M. Looseley, and B. C. Meyers. 2004. Effects fo gene expression on molecular evolution in *Arabidopsis thaliana* and *Arabidopsis lyrata*. *Mol. Biol. Evol.* 21: 1719–1726. [18]
- Yang, J., A. I. Su, and W.-H. Li. 2005. Gene expression evolves faster in narrowly than in broadly expressed mammalian genes. *Mol. Bio. Evol.* 22: 2113–2118. [18]
- Yang, Z., R. Nielsen, N. Goldman, and A.-M. K. Pedersen. 2000. Codon-substitution models for heterogeneous selection pressures at amino acid sites. *Genetics* 155: 431–449. [8]
- Yeung, K. Y., C. S. Fraley, A. Murua, A. E. Raftery, and W. L. Ruzzo. 2001. Model based clustering and data transformations for gene expression data. *Bioinformatics* 17:10:977–987. [18]
- Yoo, B. H. 1980a. Long-term selection for a quantitative character in large replicate popouations of *Drosophila melanogaster*. I: Response to selection. *Genet. Res.* 35: 1-17. [11]
- Yoo, B. H. 1980b. Long-term selection for a quantitative character in large replicate popouations of *Drosophila melanogaster*. II: Lethals and visible mutants with large effects. *Genet. Res.* 35: 19-31. [11]
- Young, S. S. Y. and H. Weiler. 1960. Selection for two correlated traits by independent culling levels. *J. Genetics* 57: 329–338. [13]
- Zeng, Z.-B. 1988. Long-term correlated response, inter-population covariation, and interspecific allometry. *Evolution* 42: 363-374. [15]